

Microlepidoptera Pannoniae meridionalis, VIII. Data to knowledge of micro-moths from Dombóvár (SW Hungary) (Lepidoptera)

IMRE FAZEKAS¹ & ARNOLD SCHREURS²

¹Biology Dept. of Regiograf Institute, Majális tér 17/A, H-7300 Komló, Hungary,
e-mail: fazekas.i@hu.inter.net;

²Conventuelenstr. 3, NL-6467 AT Kerkrade, Netherlands, e-mail: aepschreurs@hetnet.nl

FAZEKAS, I. & SCHREURS, A.: *Microlepidoptera Pannoniae meridionalis, VIII. Data to knowledge of micro-moths from Dombóvár (SW Hungary) (Lepidoptera)*.

Abstract A list of 436 species of micro-moth recorded from the area around Dombóvár-Gunaras (Tolna County, SW Hungary) is presented. Most of the material is from light traps, and several rarities have been found personally by Arnold Schreurs and Imre Fazekas between 1985 and 2009. Faunistic and biological notes on 35 species are given. Structure of genitalia and morphological characteristics of wings are illustrated with figures. Specimens are deposited in the private collections of Arnold Schreurs (NL), Willy Biesenbaum (D) and in the collection of Regiograf Institute (H). One species, *Coleophora alnifoliae* Barasch, 1934 is new to the fauna of Hungary. 14 species of Microlepidoptera are recorded as new to the fauna of the Transdanubian Hills: *Borkhausenia fuscescens* (Haworth, 1828) (Oecophoridae), *Coleophora pseudorepentis* Toll, 1960 (Coleophoridae), *C. artemisicolella* Bruand, 1855 (Coleophoridae), *C. onobrychiella* Zeller, 1849 (Coleophoridae), *Sorhagenia lophyrella* (Douglas, 1846) (Cosmopterigidae), *Aristotelia subdecurtella* (Stainton, 1858) (Gelechiidae), *Caryocolum blandulella* (Tutt, 1887) (Gelechiidae), *Gynnidomorpha alismana* (Ragonot, 1883) (Tortricidae), *Apotomis betuletana* (Haworth, 1811) (Tortricidae), *Eucosma flavispectula* Kuznetzov, 1964 (Tortricidae), *Pammene regiana regiana* (Zeller, 1849) (Tortricidae), *P. aurita* Razowski, 1992 (Tortricidae), *Phycitodes inquinatella exustella* (Ragonot, 1888) (Pyralidae) and *Catoptria permutteratella* (Herrich-Schäffer, 1848) (Crambidae).

Keywords – Microlepidoptera, faunistic, new distribution data, biology, Hungary.

Introduction

It would be hard to find another wonderful land such as the Tolna County, where continental and submediterranean characteristics are unified in a harmony. According to PAP (2007) Tolna is one of the most varied and controversial counties in Hungary. Evaluations of the country are highly contrastive. The duality of poverty and economic power, backwardness and a rich cultural heritage, easily accessible, pleasant settlements as well as increasing migration, availability of resources and favourable local conditions versus a lack of their utilisation are all characteristic features of Tolna.

Dombóvár, the second largest town in Tolna County, is 29 km from Kaposvár and 45 km from Pécs on the boundaries of three counties, Tolna, Somogy and Baranya. Gunaras, a town which is famous for its spa, is located 5 km from Dombóvár; the thermal waters

were discovered here in the 1960's, and the first bathing facilities were built in the 1980's. Nowadays it is not only a comfortable spa, but also the only rehabilitation centre in the region. The water contains fluoride and alkali-hydrogen-carbonic, thus it is mostly recommended for those who have some kind of motor disease, stomach or enteric problems or some illness of the mouth, heart or blood-system.

Scientists from Hungary and other European countries noticed the special biogeographical features of the Tolna County from as early as the 19th century. In the 20th century, famous botanists and zoologists travelled through the region. A series of smaller and larger zoological publications were published in Hungarian, Austrian and German scientific reviews. As a result, the name of Tolna County became generally known in Europe.

However, there was no systematic research undertaken in this geographical area until now. The history of lepidopterological investigation of the Tolna County goes back to the collecting activity of PILICH (1914) of Simontornya. In accordance with the usual practice of that period, he unfortunately failed to label his material, and the nomenclature he applied gave rise to several misinterpretations. More recently, additional data on micro-moths and butterflies of the area have appeared in other short publications (FARKAS 1992; FAZEKAS 1992, 2001b, 2008b, PILICH 1914).

This study presents a list of 436 Microlepidoptera species recorded from the area around Dombóvár–Gunaras (SW Hungary, Tolna County; see Fig. 1.). The account is based on material collected mainly by Arnold Schreurs (NL-Kerkade) and Willy Biesenbaum (D-Velbert-Langenberg), which is deposited in Regiograf Institute, H-Komló, and on the above-mentioned literature sources. The bulk of the information available is the result of research by Arnold Schreurs. The arrangement of the species is based on the classification still followed in Hungarian Microlepidoptera literature (FAZEKAS 2002, PASTORÁLIS 2010).

Significant species from the territory from a faunistic point of view

Abbreviation in the text: HNHM= Hungarian Natural History Museum (Budapest), JPM= Janus Pannonius Museum (Pécs).

Ornixola caudulatella (Zeller, 1839) (Gracillariidae) – Material examined: Dombóvár, Gunaras, 1♀, 12-23.07. 1999; 1♂, 15.06.2003; 2♀, 13-23.06.2006. Very local and rare on the Southern-Transdanubia: Diósviszló, Fonyód, Gyűrűfű, Kaposvár and Pécs (FAZEKAS 2001, 2002; SZEÖKE 2009). Larva oligophagous. Foodplants in Hungary: *Salix* spp.. The moth is bivoltine and flies May-June and July-August. Habitats: willow and birch mire woodlands, riverine willow-poplar woodlands, riverine ash-alder woodlands. Generally rare and local in Hungary.

Elachista alpinella Stainton, 1854 (Elachistidae) – Material examined: Dombóvár, Gunaras, 2♂, 1-15.08.1997; 2♂, 25-31.08.1998; 1♂, 26.07-7.08. 2008, det. W. Biesenbaum. Rare species with very isolated populations in Hungary. The data in the literature are vague. Sporadic records of distribution from Hungary: – “Budapest, 1912.V.13., leg. Uhrik”, in coll., HNHM (Szöcs 1973); “Pécs, 1955.V.31., leg. Nattán” (FAZEKAS 2002; SZABÓKY 1983: in coll. Nattán, JPM); Agárd, Dinnyés (PETRICH 2001). There are no verified records of *E. alpinella* from the Great Hungarian Plain and it is not recorded from western Hungary. Widespread in the western and northern parts of Europe. Larva oligophagous on *Carex* spp., and overwinters in the mine. Flight period



Fig. 1: Approximata position of the observation site of Microlepidoptera in Hungary (a)
Habitats of micro-moths at Gunaras near Dombóvár (b)




Fig. 2: *Scythris sinensis*: adult (a), underside (b), male genitalia (c), abdominal tergites of female (d-e), abdominal tergites of male (f-g). – *Scythris lampyrella*: abdominal tergites of male (h). – Distribution of *Scythris sinensis* in Hungary (i)

in Hungary unknown; Elsewhere in Europe, the moth flies from May to September, and is probably univoltine.

Scythris sinensis Felder & Rogenhofer, 1875 (Scythrididae) – According to FAZEKAS (2008b) the specimen studied is deposited in the collection of Arnold Schreurs (Netherlands): Hungary, Dombóvár, Gunaras, 10.06.2006, leg. A. Schreurs; N 46°23'49.95", E 18°10'49.11"; 10 x 10 km UTM grid code: BS 84; altitude 116 m. The Dombóvár (Gunaras) locality lies a distance of more than 150 km from the old localities in Central Hungary (Tápióság). The habitat an old, abandoned rail track in agricultural country in which there are some industrial areas. Typical habitats of the species in Hungary are places with industrial, commercial and agricultural ruderal sites. First recorded from Central Hungary (Tápióság): this specimen was found in a house; it is sooty-black, lacking the yellow spot on the forewing. Altogether, two specimens are now known from Hungary. The larva feeds on *Chenopodium album*. Moths have been collected in May and July. It is known to occur in Great Britain, Germany, Estonia, Latvia, Lithuania, Belorussia, Ukraine and Moldova and outside Europe in C. Russia, S. Siberia, E. Asia, Japan, Korea and Taiwan. *Scythris* Hübner, 1825, the largest genus in the family Scythrididae, contains more than 30 species in Hungary.

Borkhausenia fuscescens (Haworth, 1828) (Oecophoridae) – Material examined: Dombóvár, Gunaras, 2♂, 14-28.07.2007. New to the fauna of the Transdanubian Hills. According to GOZMÁNY (1958) only collected in western Hungary areas (Transdanubia). Known also from the sand region of the Kiskunság: two specimens taken at light in Ócsa (GOZMÁNY, SZABÓKY 1986). The larvae feed on a wide range of dried plant matter such as dead leaves and birds' nests, normally within a silken tube (KIMBER 2010). Moth collected in July and August.

Epicallima bruandella (Ragonot, 1889) (Oecophoridae) – Material examined: Dombóvár, Gunaras, 1♀, 1-18.08.1992. – The species has been reported in Southern Transdanubia from only two localities (FAZEKAS 2002): Baranya County, Villányi-Hills (Szászomlyó). The species occurs in white oak scrub woodlands and calcareous open rock grasslands (*Sedo sopianae–Festucetum dalmaticae*); Kaposvár. Sporadically distributed in Hungary: Budapest, Kaposvár, Parád (GOZMÁNY 1958), Jászberény, Nagykáta, Jászfelsőszentgyörgy [open sand steppe oak woodlands] (BUSCHMANN 2003), Gánt (TAKÁCS 2009). Biology: Larvae live on rotting wood particularly oak from decaying wood collected in April he raised in July several specimens. Moths have been collected in July and August in Hungary.

Batrachedra praeangusta (Haworth, 1828) (Batrachedridae) – Material examined: Dombóvár, Gunaras, 1♂, 14-28.07.2007. Distributed in Hungary: Kaposvár, Budapest, Ágasegyháza (poplar-juniper steppe woodlands), Dunaújváros (Pastorális pers.com.) and North Hungarian Mountains. Larva feeds on *Populus alba* L. and *P. tremula* L. (GOZMÁNY 1958). Adults are found in June-July.

Augasma aeratella (Zeller, 1839) (Coleophoridae) – Material examined: Dombóvár, Gunaras, 5♂, 3♀, 1-18.08.1998, det. H. v. d. Wolf. Localities in Southern Transdanubia: Kaposvár, Pécs. Sporadic and local in the Great Hungarian Plain (Szöcs 1977). The larva lives on *Polygonum aviculare* L. from October to April and the moth flies from May to August. Habitat: xerothermophilous species, found mainly in the closed loess and sand steppes, saline pasture, edge of agricultural land.

Coleophora alnifoliae Barasch, 1934 (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♂, 13-23.06.2006; 1♂, 14-28.07.2007, det. H. v. d. Wolf. According to FAZEKAS (2010), new to the fauna of Hungary. This species has not been reported before from Hungary, which is surprising, because it is widespread all over Europe wherever *Alnus* grows. It is difficult to distinguish adults of *Coleophora alnifoliae* from *C. milvi-*




Fig. 3: *Coleophora alnifoliae*: adult (a), sack (b), female genitalia (c)

pennis Zeller, 1839, *C. limosipennella* (Duponchel, 1843) and *C. badiipennella* (Duponchel, 1843). Also the male genitalia are difficult to identify. The female genitalia are easier. Distribution: Europe, Caucasus, Central Siberia, Canada and U.S.A. (BALDIZZONE et al. 2006). According to BIESENBAUM and WOLF (1999) "Die Raupen fertigen zuerst in einem Jugendsack, später in einem Blattsack und minieren an Erle (*Alnus*)". Adult's univoltine between June and August.

Coleophora trifolii (Curtis, 1832) (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♀, 7-15.08.2000; 1♀, 14-28.07.2007; 4♂, 4♀, 14-24.07.2004, det. H. v. d. Wolf. Localities in Southern Transdanubia: Kárász, Komló-Mecsekjánosi, absent in Somogy County. Larvae known on *Melilotus* spp. and the moth flies from June to August. Sporadic and distributed in meadow and grassland habitat in Hungary, mostly in the Hungarian Plain and very local in mountainous areas (Bükk Mts. and Mecsek Mts.): Ágasegyháza, Győr, Fülpöháza, Izsák, Jászberény, Kárász, Komló, Kunadacs, Kunszentmiklós, Miskolc, Nagyvisnyó. Known also from the Bátorliget Nature Conservation Areas (ÁCS et al 1990). The Bátorliget Nature Conservation Area is one of the oldest protected territories of Hungary. Distribution: in Palaearctic and adventives from Canada to U.S.A.

Coleophora onobrychiella Zeller, 1849 (= *arenariella* Zeller, 1865) (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♂, 1-18.08.1998; 1♀, 5-15.06.2003, det. H. v. d. Wolf. – New to the fauna of the Transdanubian Hills. According to GOZMÁNY (1965), larvae feed on *Astragalus arenarius* L. in Central and North Europe. In recent times known on *Astragalus onobrychis* L. (Pastorális pers. com.; from Örkény, in coll.




Fig. 4: Distribution map of *Coleophora trifolii* in Hungary

Ivan Richter). The foodplant is unknown in Hungary. Biology in Hungary: no data except for collection dates of moth; June and August. The distribution in Hungary: Velencei Hills. and Budai Mts. (Transdanubian Mountains), Csákberény and Örkény (Pastorális pers. com). Distribution: Widely distributed in Palaearctic region from France to China.

Coleophora pseudorepentis Toll, 1960 (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♂, 1-18.08.1992; 1♂, 1-18.08.1998, det. H. v. d. Wolf. New to the fauna of the Transdanubian Hills. Very rare and local in Hungary: Bükk Mts. and Vértes Mts. Known in Europe from Southern Ural to France.

Coleophora artemisicolella Bruand, 1855 (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♂, 1-15.08.1997; 2♂, 3♀, 1-18.08.1998; 3♂, 2♀, 28.07-08.08.2008, det. H. v. d. Wolf. – New to the fauna of the Transdanubian Hills. It is a Transpalaearctic species. Larva monophagous on *Artemisia vulgaris* L. and the adults fly from June to August. The distribution in Hungary: Bátorliget, Budapest, Ócsa (GOZMÁNY 1956), Törökbalint, Érd-Elvira (SZIRÁKI 1980). The majority of those collected by Sziráki were recovered from “Atralin” traps (pheromone preparations produced in Roumania). Type synthetic attractant: sex pheromone for *Anarsia lineatella* Zeller, 1839 (Gelechiidae).

Coleophora chrysanthemi Hofmann, 1869 (Coleophoridae) – Material examined: 1♂, “Tolna Komitat, Dombóvár, Gunarasfürdő, UTM Cood: BS 84, 1-10.VIII.1997 leg. et coll. A. Schreurs, det. H. v. Wolf. According to FAZEKAS (2001b) “Habitat: Die Fundorte in Mittel- und Südeuropa liegen in Hügel- und Berggebieten. Das Areal liegt in den Zonen der Laubwälder des gemäßigten Klimas. Es erreicht im Süden (Italien) nicht den Gürtel der mediterranen Hartlaubwälder, im Norden aber (Finnland) fließt er in die Taigawaldzonen ein. In Ungarn wurde die Art in einer unter submediterranem Einfluss stehenden lößhaltigen kultivierten Flur gefunden (SW-Ungarn, Dombóvár, Gunarasfürdő,




Fig. 5: Distribution map of *Coleophora striatipennella* in Hungary

ca. 120 m, UTM Coord: BS 84). Im Gebiet herrschten früher *Fraxino pannonicae-Ulmetum* (CORINE Coord: 44.431) beziehungsweise *Orno-Quercetum* (CORINE Coord: 41.73744) Waldassoziationen vor. Heute ist das Gebiet durch Ackenwirtschaft und Erholungstourismus gekennzeichnet.” – New material examined: 2♂, 1-18.08.1998, det. H. v. d. Wolf. Larva monophagous on *Chrysanthemum corymbosum* L. and the moths fly from early May to August. European chorotype, found sporadically distributed from Italy to Central Europe and Finland.

Coleophora striatipennella Nylander, 1848 (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♀, 1-18.08.1998, 1♂, 12-24.07.2004, det. H. v. Wolf. Localities in Southern Transdanubia: Kaposvár, Simontornya, but unknown in Mecsek Mountains and Villányi Hills. Known also from the sand region of the Kiskunság: two specimens taken at light in Ócsa (GOZMÁNY, SZABÓKY 1986). The landscape conservation area at Ócsa, south-east of Budapest, was proposed as worthy of protection by lepidopterists and subsequently it was soon established as a protected dual habitat of marshy alder woods and peat meadows in the first half of the fifties. The Kiskunság region is also characterised by the great specific richness of the submediterranean psammophilous fauna and locally in habitats of high humidity. Larva polyphagous on *Cerastium*, *Moehringia*, *Stellaria* and *Pulicaria* spp. The moth flies from June to July (GOZMÁNY 1956). New the period of flight from August. *C. striatipennella* is a meso- and semi hygrophilous species in Hungary. It is a Transpalaearctic species.

Coleophora adspersella Bernander, 1939 (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♂, 1-15.08.1992; 1♂, 2♀, 12.24.07.2004; 1♀, 28.07.-8.08.2004, det. H. v. Wolf. The distribution in Hungary: Budapest, Velencei-tó (Gozmány 1965). Larva polyphagous on *Atriplex*, *Chenopodium* and *Juncus* spp. The moth flies from June to August. Known from Japan to Europe, but unknown in Asia Minor and North Africa.

Coleophora bernoulliella (Goeze, 1783) (Coleophoridae) – Material examined: Dombóvár, Gunaras, 3♂, 2♀, 5-15.06.2003; 1♂, 1♀, 13-23.06.2006, det. H. v. Wolf. – According to BALDIZZONE et al. (2006) *Tinea anatipennella* Hübner, 1796 corrected original spelling for *Tinea anatipenella* (ICZN 1999: Art. 19.2 & 32.2.2); emendation proposed by STEPHENS (1929: (II) 211)].

Coleophora caespitiella Zeller, 1839 (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♂, 5-15.06.2003, det. H. v. Wolf; 1♂, 8.06.2009, det. Fazekas. It is known from only six places in Hungary: Büksérc, Kaposvár, Miskolc, Ócsa, Szakonyfalu. The larva lives on *Juncus* spp. The moth flies from May to June. The species is distributed from Iran, Caucasus, Southern Ural and Asia Minor to Europe.

Coleophora trifariella Zeller, 1849 (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♀, 5-15.06.2003, det. H. v. Wolf. Localities in Southern Transdanubia: Kaposvár. The first specimen of *C. trifariella* in Hungary was caught in a continuously working light-trap near Kaposvár 1962. Larva polyphagous on *Cytisus*, *Genista* and *Sarrothamnus* spp. The moth flies from June to July (GOZMÁNY 1956, SZÖCS 1977). A larva starts to feed in autumn and has left its mine by the spring. *C. trifariella* occurs almost throughout Southern and Central Europe.

Coleophora motacillella Zeller, 1849 (= *palumbipennella* Toll, 1952; *szekessyi* Gozmány, 1956) (Coleophoridae) – Material examined: Dombóvár, Gunaras, 1♂, 1-15.08.1991, det. and coll. H. v. Wolf. Localities in southern Transdanubia: Kaposvár, Rinyatamási. Records from Hungary are unconfirmed and doubtful. According to GOZMÁNY (1956), the larva lives on *Atriplex* and *Chenopodium* spp. Adults fly from June to early September. Occurs from Altai region to Central Europe and France.

Mompha epilobiella ([Denis & Schiffermüller], 1775) (Momphidae) – Material examined: Dombóvár, Gunaras, 1♀, 14-28.07.2007, det. J. C. Koster. In July and August a




Fig. 6: Male (a) and female genitalia (b) of *Mompha epilobiella* (Koster & Sinev 2003)

very common species in Palaearctic. Distribution in Europe to southern Scandinavia in the north and Asia Minor, the Caucasus, submontane regions of Central Asia (KOSTER & SINEV 2003). Sporadic and very local in Hungarian mountainous areas (GOZMÁNY 1958: Bükk Mts.)

Sorhagenia lophyrella (Douglas, 1846) (Cosmopterigidae) – Material examined: Dombóvár, Gunaras, 1♀, 13-23.06.2006, det. J. C. Koster. – New to the fauna of the Transdanubian Hills, and this is the first record known from this area. Very similar species in Hungary: *S. rhamniella* (Zeller, 1839) and *S. janiszewskae* Riedl, 1962. According to KOSTER & SINEV (2003), *S. lophyrella* differs from *S. rhamniella* in the generally some what lighter forewings and paler grey hindwings, but it can only distinguished with certainty by the genitalia. It is local in Hungary: Bakony Mts., Budapest, Fót. The literary data are vague (see SZÖCS 1973). Mesophilous species, probably univoltine, adults fly from early May to mid-August. The species feeds on buckthorn (*Rhamnus cathartica* L.) and *R. saxatilis* L., at first in the bud, then later between spun leaves. Occurs from Caucasus region, Asia Minor to Europe.

Caryocolum blandulella (Tutt, 1887) (Gelechiidae) – Material examined: Dombóvár, Gunaras, 1♂, 18-24.07.1994; 1♂, 1-15.08.1996; 1♂, 28.07.-8.08.2008, det. O. Karsholt. New to the fauna of the Transdanubian Hills. There is only one reliable reference from the area of Hungary from June 1988 when Cs. Szabóky caught a specimen in Budapest (Sas-hegy). The voucher specimen is in the collection of Cs. Szabóky (Budapest). *C. blandulella* is apparently very rare and local in Hungary, but could be overlooked and therefore careful search should be made. According to ELSNER et al. (1999), known in sandy habitats. Habitat in Hungary: colline dry degraded grasslands, semi-natural road verges, semi-natural vegetation of abandoned fields (Dombóvár region) and rock steppes (Budapest).

Aristotelia subdecurtella (Stainton, 1858) (Gelechiidae) – Material examined: Dombóvár, Gunaras, 1♀, 05.15.06.2003, det. O. Karsholt, in coll. J. B. Wolschrijn (NL). New to the fauna of the Transdanubian Hills. Local in Central Europe. Occurrence in Hungary: Farmos, Jászberény, Pécsely, Pomáz, Terecseny (BUSCHMANN 2003, SZABÓKY 1994, 2009). Vertical distribution: in Hungary found up to about 300 m. Adults occur from early June to late August. Larvae polyphagous on species of *Lythrum*, *Stachys* and *Veronica*.

Chrysoesthia sexguttella (Thunberg, 1794) (Gelechiidae) – Material examined: Dombóvár, Gunaras, 1♀, 20.08.2009, gen. prep. Schreurs, det. Fazekas. Only known from one locality in Transdanubian Hills (Kaposvár), 56 years before (FAZEKAS 2001). The absence of records from Mecsek Mountains and Villányi Hills remains obscure (Fazekas 2002, 2007). Sporadic and local in Transdanubian Mountains and north Hungarian Mountains, flying in two generation from May to September. The larva has been recorded on various Chenopodiaceae and Amaranthaceae species. Holarctic: widely distributed from Ireland to Japan and Canada. Similar species: *Ch. verrucosa* Tokár, 1999.

Syncopacma ochrofasciella (Toll, 1936) (Gelechiidae) – Material examined: Dombóvár, Gunaras, 2♂, 25-31.08.1998; 1♂, 14-28.07.2007, gen. prep. Schreurs, Nr. 982. There are only very old records from Hungary: Isaszeg, Kaposvár, Szár (GOZMÁNY 1958). We had to wait for the discovery of a new population of *S. ochrofasciella* until 1998, when A. Schreurs noticed the imagos of the moth during a light trap survey in Gunaras. The larva lives on *Astragalus glycyphyllos* L. Bivoltine, adults in May and August. Chorotype: Eurosiberian. Very local in Central Europe.

Gynnidomorpha alismana (Ragonot, 1883) (Tortricidae) – Material examined: Dombóvár, Gunaras, 1♂, 1♀, 12-24.07.2004, , det. H. v. Wolf. New to the fauna of the




Fig. 7: Adult and distribution map of *Eucosma flavispectula* in Hungary

Transdanubian Hills. There is only one reliable record from this area of Hungary, in July 1929 when Ludwig Ostheder caught a specimen in Pusztapeszér (FAZEKAS 1994b: Figs. 2. and 3). The voucher specimen is in the collection of Zoologische Staatssammlung of Munich; gen. prep. I. Fazekas, No. 2746.




Fig. 8: Adults and copulatory organs of *Catoptria permutterella*: – adults; (a) Telkibánya, (b) Istvánkút: – female genitalia: (c) Dombóvár (gen. prep. Schreurs): – male genitalia (valva); (d) Sopron, (e) Istvánkút, (f) Szakonyfalu, (g) Telkibánya, (h) Istvánkút, (i) Telkibánya (del. et gen. prep. Fazekas)

Cochylis flavigiliana (Westwood, 1854) (Tortricidae) – Material examined: Dombóvár, Gunaras, 2♂, 1♀, 1-15.08.1991, det. I. Fazekas, gen. prep. No. 2588, 2589; 2♂, 14-28.07.2007; 2♂, 1♀, 1-19.08.1991, leg. et coll. R. Beyer (D-Linnich), det. I. Fazekas (see FAZEKAS 1994a). The species is local in southwest Hungary: Mecsek Mountains and the Dombóvár region. Known only in the summer generation in some areas. Biology: oligophagous; recorded foodplants are *Scabiosa* spp. and *Knautia arvensis* (L.) Coul. Typical habitat in Hungary: moist rich fens, eu- and mesotrophic meadows, colline and montane hay meadows and acid grasslands. Chorotype: Temperate-Meridional; West-mid-Palaearctic.

Apotomis betuletana (Haworth, 1811) (Tortricidae) – Material examined: Dombóvár, Gunaras, 2♂, 26.07.-8.08.2008. New to the fauna of the Transdanubian Hills. There is only one record known from this area. Occurrence in Hungary: Nyírád, Nagyvisnyó (Bükk Mts.), Parád, Sástó (Mátra Mts.).

Eucosma flavispectula Kuznetzov, 1964 (Tortricidae) – Material examined: Dombóvár, Gunaras, 1♂, 1♀, 1-15.08.1991; 1♀, 1-18.08.1992; 2♀, 1-15.08.1997; 1♂, 1-15.08.1998; 4♂, 2♀, 12-23.07.1999; 3♂, 1♀, 12-24.07.2004; 2♂, 1♀, 7-15.08.2000. – New to the fauna of the Transdanubian Hills. Sporadic and very local in Hungary: Bélmegyer, Csikóspuszta (Körös-Maros Nemzeti Park), Fülöpháza (Kiskunság), Kunpeszér, Salföld (Ábrahám-hegy), Agárd. According to SZABÓKY (2004) the larva feeds on *Centaurea pannonica* (Heuff.), besides on *C. jacea* L. (RAZOWSKI 2003). The moth flies from June to August in Hungary. Distribution in Palaearctic: Russian Far East, Mongolia, South Siberia, Kazakhstan, western and central parts of East Europe (RAZOWSKI 2003).

Pammene regiana regiana (Zeller, 1849) (Tortricidae) – Material examined: Dombóvár, Gunaras, 2♀, 13.23.06.2006. New to the fauna of the Transdanubian Hills. Obscure occurrence in Hungary. Very old references from GOZMÁNY (1968). Life history: larva from August and September to April, in seeds of species of maple (*Acer platanoides* L., *A. pseudoplatanus* L., *A. campestre* L.). The moth flies in one generation from May to June. Distribution: *P. regiana regiana* subspecies distributed from east Europe to Scandinavia and Scotland (RAZOWSKI 2003).

Pammene aurita Razowski, 1992 (Tortricidae) – Material examined: Dombóvár, Gunaras, 1♀, 18-28.07.1994; 1♀, 1-15.08.1996; 1♀, 7-15.08.2000; 1♂, 2♀, 14-28.08.2009. New to the fauna of the Transdanubian Hills. Unknown elsewhere in Hungary. Very old references from GOZMÁNY (1968). The larvae, in August and September, feed inside the seeds of *Acer pseudoplatanus* L., and their presence is sometimes betrayed by small exit holes in the seeds themselves. The moth flies from July to August. Distribution in Europe: Till now known from east Ukraine to France and British Is.

Grapholita discretana (Wocke, 1861) (Tortricidae) – Material examined: Dombóvár, Gunaras, 1♀, 12-23.07.1999. According to PASTORALIS (2010), the first record in Hungary originated from GOZMÁNY (1968). Very local in Transdanubian Hills: Kaposvár (SZABÓKY 1983), Mecsek Mts. (FAZEKAS 2002). The moth flies from May to July, probably in single generation yearly. Widely distributed from Kazakhstan to Europe. Larva from July to September, and after hibernation in April, on *Humulus lupulus* L.

Phycitodes inquinatella exustella (Ragonot, 1888) (Pyralidae) – Material examined: Dombóvár, Gunaras, 1♂, 1♀, 1-15.08.1998; 2♂, 12-13.07.1999, det. J. Asselsberg. New to the fauna of the Transdanubian Hills. Occurrence in Hungary: Bakony- and Bükk Mts. (FAZEKAS 1993, 1996). Limited information available about habitat preference (FAZEKAS 1996): acid open rock grasslands (Tihany), slope steppes (Bükk Mts.), from 200 m up to 400 m above sea-level. Probably univoltine in Hungary, in July and August. Male genitalia and forewing drawings: FAZEKAS (1996: Fig. 1. a-b.). The subspecies is distributed

in Europe from Hungary through Austria and Germany to France. Replaced by allopatric sister subspecies in Palaearctic: nominate subspecies known in Asia Minor, Crete, Cyprus, Balkans and Roumania; subspecies *ravonella* Pierce, 1937 occurs in Italy, Sicily, Corsica; subspecies *canariella* Rebel, 1892, flies in south Spain, north Africa and Canary Islands.

Phycitodes albatella pseudonimbella Benetinck, 1937 (Pyralidae) – Material examined: Dombóvár, Gunaras, 1♂, 14-28.08.2009. According to FAZEKAS (1998): "Die Unterart ist in Ungarn lokal und nicht weitverbreitet. Ihre Habitate sind von trocknen, tiefebenischen Wiesen bis zu dunstigen Talgebiete de Mittelgebirgen zu finden." Occurrence in Hungary: Bakony Mts., Bükk Mts., Mátra Mts., Mecsek Mts. and the Great Hungarian Plain (FAZEKAS 1998).

Catoptria permutterella (Herrich-Schäffer, 1848) (Crambidae) – Material examined: Dombóvár, Gunaras, 14-18.07.2007, gen. prep. Schreurs, No. 957. New to the fauna of the Transdanubian Hills. Occurrence in Hungary (FAZEKAS 1986: Abb. 9.): Sopron–Bánfalva, Szakonyfalu, “Zempléner Gebirge, Istvánkút”, Telkibánya, Jósvafő. Restricted to the lowlands from 120 m up to about 500 m above sea-level. Life history: larva lives on moss species and hibernates; adults fly in July and August. A very common species in west and north Hungary. The vernal generation is unknown in Hungary. Costal arm of valva in male genitalia are rather variable (see FAZEKAS 1986: Abb. 1-6.)

List of species

Comment: A note was made of the species labelled with the star.

NEPTICULIDAE

- Stigmella aceris* (Frey, 1857)
Stigmella plagicolella (Stainton, 1854)
Ectoedemia louisella (Sircom, 1847)

OPOSTEGIDAE

- Opostega spatulella* (Herrich-Schäffer, 1855)
Pseudopostega auritella (Hübner, 1813)
Pseudopostega crepusculella (Zeller, 1839)

TISCHERII

- Tischeria ekebladella* (Bjerkander, 1795)
Coptotriche marginea (Haworth, 1828)

Monopis mo

- ## GRACILLARIIDAE

Parectopa robinella

- Caloptilia roscipennella* (Hubner, 1796)
Caloptilia stigmatella (Fabricius, 1781)
Calybites phasianipennella (Hübner, 1813)
Aspilapteryx tringipennella (Zeller, 1839)
Eucalybites auroguttella (Stephens, 1835)
Ornixola caudulatella (Zeller, 1839)*

Parornix anglicella (Stainton, 1850)

- Paromix finimitella* (Zeller, 1850)
Cameraria ohridella Deschka & Dimić, 1986
Phyllonorycter robiniella (Clemens, 1859)
Phyllonorycter abrasella (Duponchel, 1843)
Phyllonorycter manni (Zeller, 1846)
Phyllonorycter maestingella (Müller, 1764)
Phyllonorycter tenerella (Joannis, 1915)
Phyllonorycter ulmifoliella (Hübner, 1817)
Phyllonorycter lantanella (Schrank, 1802)
Phyllonorycter spinicolella (Zeller, 1846)
Phyllonorycter blancardella (Fabricius, 1781)
Phyllonorycter oxyacanthae (Frey, 1856)
Phyllonorycter populifoliella (Treitschke, 1833)
Phyllonorycter pastorella (Zeller, 1846)
Phyllonorycter comparella (Duponchel, 1843)
Phyllonorycter medicaginella (Gerasimov, 1930)
Phyllocnistis saligna (Zeller, 1839)
Phyllocnistis xenia Hering, 1936
Phyllocnistis unipunctella (Stephens, 1834)

YPONOMEUTIDAE

- Scythropia crataegella* (Linnaeus, 1767)
Yponomeuta evonymella (Linnaeus, 1758)
Yponomeuta plumbella ([Denis & Schiffermüller], 1775)
Yponomeuta sedella Treitschke, 1832

- Swammerdamia pyrella* (de Villers, 1789)
Argyresthia brockeella (Hübner, 1813)
Argyresthia goedartella (Linnaeus, 1758)
Argyresthia conjugella (Zeller, 1839)
Argyresthia pruniella (Clerck, 1759)
- YPSOLOPHIDAE**
Ypsolopha ustella (Clerck, 1759)
- PUTELLIDAE**
Plutella xylostella (Linnaeus, 1758)
- GLYPHIPTERIGIDAE**
Orthotelia sparganella (Thunberg, 1788)
Glyptipterix simpliciella (Stephens, 1834)
- BEDELLIIDAE**
Bedellia somnulentella (Zeller, 1847)
- LYONETIIDAE**
Leucoptera sinuella (Reutti, 1853)
Lyoneta clerkella (Linnaeus, 1758)
- ETHMIIDAE**
Ethmia quadrillella (Goeze, 1783)
Ethmia bipunctella (Fabricius, 1775)
Ethmia haemorrhoidella Eversmann, 1844
- DEPRESSARIIDAE**
Agonopterix alstroemeriana (Clerck, 1759)
Agonopterix heracliana (Linnaeus, 1758)
Depressaria depressana (Fabricius, 1775)
Depressaria pulcherrimella Stainton, 1849
- ELACHISTIDAE**
Elachista utorella Frey, 1856
Elachista alpinella Stainton, 1854*
Elachista maculicerusella Bruand, 1859
Elachista argentella (Clerck, 1759)
Elachista bisulcella (Duponchel, 1843)
- SCYTHRIDIDAE**
Scythris limbella (Fabricius, 1775)
Scythris sinensis Felder & Rogenhofer, 1875*
Parascythris muelleri (Mann, 1871)
- OECOPHORIDAE**
Bisigna procerella ([Denis & Schiffermüller]), 1775
Metralampa cinnamomea (Zeller, 1839)
Hofmannophila pseudospretella (Stainton, 1849)
Borkhausenia fuscescens (Haworth, 1828)*
Crassa tinctella (Hübner, 1796)
Crassa unitella (Hübner, 1796)
Epicallima bruandella (Ragonot, 1889)*
Oecophora bractella (Linnaeus, 1758)
- Alabonia staintoniella* (Zeller, 1850)
Harpella forficella (Scopoli, 1763)
Carcina quercana (Fabricius, 1775)
- BATRACHEDRIDAE**
Batrachedra paeangusta (Haworth, 1828)*
Batrachedra pinicolella (Zeller, 1839)
- COLEOPHORIDAE**
Augasma aeratella (Zeller, 1839)*
Coleophora alnifoliae Barasch, 1934*
Coleophora limosipennella (Duponchel, 1845)
Coleophora prunifoliae Doets, 1944
Coleophora trifolii (Curtis, 1832)*
Coleophora lusciniaeepennella (Treitschke, 1833)
Coleophora frischella (Linnaeus, 1758)
Coleophora alcyonipennella (Kollar, 1832)
Coleophora lineolea (Haworth, 1828)
Coleophora onobrychiella Zeller, 1849*
Coleophora colutella (Fabricius, 1794)
Coleophora trifariella Zeller, 1849*
Coleophora ballotella (Fischer von Röslerstamm, 1839)
Coleophora serpylletorum Hering, 1889
Coleophora bernoulliella Goeze, 1783)*
Coleophora zelleriella Heinemann, 1854
Coleophora gallipennella (Hübner, 1796)
Coleophora conspicuella Zeller, 1849
Coleophora partitella Zeller, 1849
Coleophora adjunctella Hodgkinson, 1882
Coleophora glaucicolella Wood, 1892
Coleophora taeniipennella Herrich-Schäffer, 1855
Coleophora therinella Tengström, 1848
Coleophora saxicolella (Duponchel, 1843)
Coleophora motacillella Zeller, 1849*
Coleophora sternipennella (Zetterstedt, 1839)
Coleophora caespitiella Zeller, 1839*
Coleophora versurella Zeller, 1849
Coleophora artemisicolella Bruand, 1855*
Coleophora chrysanthemi Hofmann, 1869*
Coleophora peribenanderi Toll, 1943*
Coleophora trochilella (Duponchel, 1843)
Coleophora inulae Wocke, 1877
Coleophora striatipennella Nylander, 1848*
Coleophora argentula (Stephens, 1834)
Coleophora pseudorepentis Toll, 1960*
Coleophora follicularis (Vallot, 1802)
Coleophora adspersella Bernander, 1939*
Coleophora clypeiferella Hofmann, 1871
Coleophora binotapennella (Duponchel, 1843)
Coleophora unipunctella Zeller, 1849
- MOMPHIDAE**
Mompha ochraceella (Curtis, 1839)
Mompha epilobiella ([Denis & Schiffermüller], 1775)*

BLASTOBASIDAE

Blastobasis phycidella (Zeller, 1839)

AUTOSTICHIDAE

Oegoconia deauratella (Herrich-Schäffer, 1854)

COSMOPTERIGIDAE

Limnaecia phragmitella Stainton, 1851

Cosmopterix orichalcea Stainton, 1861

Cosmopterix scribaiella (Zeller, 1850)

Cosmopterix lienigiella Lienig & Zeller, 1846

Pyroderces argyrogrammos (Zeller, 1847)

Eteobalea serratella Treitschke, 1833

Sorhagenia lophyrella (Douglas, 1846)*

GELECHIIDAE

Megacraspedus dolosellus (Zeller, 1839)

Aristotelia decurtella (Hübner, 1818)

Aristotelia subdecurtella (Stainton, 1858)*

Chrysoesthia sexguttella (Thunberg, 1794)*

Isophrictis striatella ([Denis & Schiffermüller], 1775)

Metzneria metzneriella (Stainton, 1851)

Monochroa tenebrella (Hübner, 1817)

Monochroa lucidella (Stephens, 1834)

Monochroa palustrellus (Douglas, 1850)

Monochroa arundinetella (Boyd, 1868)

Monochroa hornigi (Staudinger, 1883)

Eulamprotes atrella ([Denis & Schiffermüller], 1775)

Bryatropa terrella ([Denis & Schiffermüller], 1775)

Bryatropa basaltinella (Zeller, 1839)

Bryatropa senectella (Zeller, 1839)

Recurvaria nanella ([Denis & Schiffermüller], 1775)

Recurvaria leucatella (Clerck, 1759)

Coleotechnites piceaella (Kearfott, 1903)

Exoteleia dodecella (Linnaeus, 1758)

Parastenolechia nigrinotella (Zeller, 1847)

Teleiodes vulgella ([Denis & Schiffermüller], 1775)

Teleiodes luculella (Hübner, 1813)

Carpotolechia proximella (Hübner, 1796)

Pseudotelphusa paripunctella (Thunberg, 1794)

Gelechia senticetella (Staudinger, 1859)

Gelechia turpella ([Denis & Schiffermüller], 1775)

Scrobipalpa acuminatella (Sircom, 1850)

Scrobipalpa proclivella (Fuchs, 1886)

Scrobipalpa atriplicella (Fischer von Röslerstamm, 1841)

Caryocolum proxima (Haworth, 1828)

Caryocolum blandulella (Tutt, 1887)*

Syncopacma cinctella (Clerck, 1759)

Syncopacma ochrofasciella (Toll, 1936)*

Aproaerema anthyllidella (Hübner, 1813)

Anacampsis populella (Clerck, 1759)

Anacampsis blattariella (Hübner, 1796)

Anacampsis timidella (Wocke, 1887)

Crossobela trinotella (Herrich-Schäffer, 1839)

Anarsia spartiella (Schrank, 1802)

Nothris verbascella ([Denis & Schiffermüller], 1775)

Dichomeris ustalella (Fabricius, 1794)

Dichomeris derasella ([Denis & Schiffermüller], 1775)

Dichomeris limosella (Schläger, 1849)

Dichomeris rasilella (Herrich-Schäffer, 1854)

Brachmia dimidiella ([Denis & Schiffermüller], 1775)

Brachmia blandella (Fabricius, 1798)

Helcystogramma triannulella (Herrich-Schäffer, 1854)

Helcystogramma rufescens (Haworth, 1828)

Sitotroga cerealella (Oliver, 1789)

LIMACODIDAE

Apoda limacodes (Hufnagel, 1766)

Heterogena asella ([Denis & Schiffermüller], 1775)

ZYGAENIDAE

Jordanita globulariae (Hübner, 1793)

Adscita statices (Linnaeus, 1758)

Zygaena purpuralis (Brünnich, 1763)

Zygaena carniolica (Scopoli, 1763)

Zygaena loti ([Denis & Schiffermüller], 1775)

Zygaena viciae ([Denis & Schiffermüller], 1775)

Zygaena filipendulae (Linnaeus, 1758)

Zygaena lonicerae (Scheven, 1777)

COSSIDAE

Cossus cossus (Linnaeus, 1758)

Zeuzera pyrina (Linnaeus, 1761)

Phragmataecia castaneae (Hübner, 1790)

TORTRICIDAE

Phalonidia manniiana (Fischer von Röslerstamm, 1839)

Phalonidia contractana (Zeller, 1847)

Gynnidomorpha luridana (Gregson, 1870)

Gynnidomorpha alismana (Ragonot, 1883)*

Agapeta hamana (Linnaeus, 1758)

Agapeta zoegana (Linnaeus, 1767)

Aethes hartmanniana (Clerck, 1759)

Aethes margaritana (Haworth, 1811)

Aethes smethmanniana (Fabricius, 1781)

Aethes tesserana ([Denis & Schiffermüller], 1775)

Aethes bilbaensis (Rössler, 1877)

Aethes cnicana (Westwood, 1854)

Aethes rubrigana (Treitschke, 1830)

Cochylidia richteriana (Fischer von Röslerstamm, 1837)

Cochylidia implicitana (Wocke, 1856)

Diceratura ostrinana (Guenée, 1845)

Cochylis nana (Haworth, 1811)

Cochylis flaviciliiana (Westwood, 1854)*

Cochylis hybridella (Hübner, 1813)

Cochylis dubitana (Hübner, 1799)

Tortrix viridana (Linnaeus, 1758)

Aleimma loeflingiana (Linnaeus, 1758)

Acleris lorquiniana (Duponchel, 1835)

Acleris variegana ([Denis & Schiffermüller], 1775)

- Acleris notana* (Donovan, 1806)
Cnephasia incertana (Treitschke, 1835)
Cnephasia pasiuana (Hübner, 1799)
Cnephasia chrysanthaea (Duponchel, 1843)
Eulia ministrana (Linnaeus, 1758)
Pseudargyrotoza conwagana (Fabricius, 1775)
Philedone gerningana ([Denis & Schiffermüller], 1775)
Archips podana (Scopoli, 1763)
Archips xylosteana (Linnaeus, 1758)
Archips rosana (Linnaeus, 1758)
Ptycholoma lecheana (Linnaeus, 1758)
Pandemis cerasana (Hübner, 1786)
Pandemis heparana ([Denis & Schiffermüller], 1775)
Pandemis dumetana (Treitschke, 1835)
Aphelia paleana (Hübner, 1793)
Dichelia histriionana (Frölich, 1828)
Clepsis spectrana (Treitschke, 1830)
Adoxophyes orana (Fischer von Röslerstamm, 1834)
Bactra furfurana (Haworth, 1811)
Bactra lancealana (Hübner, 1799)
Bactra robustana (Christoph, 1872)
Endothenia gentianaeana (Hübner, 1799)
Endothenia quadrimaculana (Haworth, 1811)
Apotomis lineana ([Denis & Schiffermüller], 1775)
Apotomis betuletana (Haworth, 1811)*
Orthotaenia undulana ([Denis & Schiffermüller], 1775)
Heda salicella (Linnaeus, 1758)
Heda nubiferana (Haworth, 1811)
Heda pruniana (Hübner, 1799)
Celypha rufana (Scopoli, 1763)
Celypha striana ([Denis & Schiffermüller], 1775)
Celypha cespitana (Hübner, 1817)
Celypha lacunana ([Denis & Schiffermüller], 1775)
Celypha rivulana (Scopoli, 1763)
Lobesia abscisana (Doubleday, 1849)
Lobesia reliquana (Hübner, 1825)
Ancylis laetana (Fabricius, 1775)
Ancylis diminutana (Haworth, 1811)
Ancylis apicella ([Denis & Schiffermüller], 1775)
Ancylis paludana (Barrett, 1871)
Ancylis badiana ([Denis & Schiffermüller], 1775)
Ancylis achatana ([Denis & Schiffermüller], 1775)
Ancylis mütterbacheriana ([Denis & Schiffermüller], 1775)
Rhopobota naevana (Hübner, 1817)
Spilonota ocellana ([Denis & Schiffermüller], 1775)
Spilonota laricana (Heinemann, 1863)
Gibberifera simplana (Fischer von Röslerstamm, 1836)
Epinotia trigonella (Linnaeus, 1758)
Epinotia abbreviana (Fabricius, 1794)
Epinotia immundana (Fischer von Röslerstamm, 1839)
Epinotia nanana (Treitschke, 1835)
Epinotia demarniana (Fischer von Röslerstamm, 1840)
Epinotia subocellana (Donovan, 1806)
Epinotia tetraquetra (Haworth, 1811)
Epinotia tenerana ([Denis & Schiffermüller], 1775)
- Epinotia tedella* (Clerck, 1759)
Epinotia bilunana (Haworth, 1811)
Epinotia nisella (Clerck, 1759)
Zeiraphera griseana (Hübner, 1799)
Zeiraphera insertana (Fabricius, 1794)
Pelochrista caecimacula (Hübner, 1799)
Eucosma cana (Haworth, 1811)
Eucosma hohenwarthiana ([Denis & Schiffermüller], 1775)
Eucosma flavispectula Kuznetsov, 1964*
Eucosma albidulana (Herrich-Schäffer, 1851)
Eucosma metzneriana (Treitschke, 1830)
Eucosma conterminana (Guenée, 1845)
Eucosma lacteana (Treitschke, 1835)
Gypsonoma minutana (Hübner, 1799)
Gypsonoma dealbana (Frölich, 1828)
Gypsonoma oppressana (Treitschke, 1835)
Gypsonoma nitidulana (Lienig & Zeller, 1846)
Epiblema scutulana ([Denis & Schiffermüller], 1775)
Epiblema foenella (Linnaeus, 1758)
Epiblema junctana (Herrich-Schäffer, 1856)
Epiblema grandaevana (Lienig & Zeller, 1846)
Notocelia cynosbatella (Linnaeus, 1758)
Notocelia uddmanniana (Linnaeus, 1758)
Notocelia roborana ([Denis & Schiffermüller], 1775)
Dichrorampha plumbana (Scopoli, 1763)
Dichrorampha acuminatana (Lienig & Zeller, 1846)
Dichrorampha simpliciana (Haworth, 1811)
Dichrorampha petiverella (Linnaeus, 1758)
Cydia nigricana (Fabricius, 1794)
Cydia oxytropidis (Martini, 1912)
Cydia succedana ([Denis & Schiffermüller], 1775)
Cydia pomonella (Linnaeus, 1758)
Cydia splendana (Hübner, 1799)
Lathronympha strigana (Fabricius, 1775)
Grapholita discretana (Wocke, 1861)*
Grapholita compositella (Fabricius, 1775)
Grapholita funebrana (Treitschke, 1835)
Grapholita tenebrosana (Duponchel, 1843)
Pammene regiana (Zeller, 1849)*
Pammene aurita Razowski, 1992*
Strophedra nitidana (Fabricius, 1794)
- CHOREUTIDAE**
Anthophila fabriciana (Linnaeus, 1767)
- EPERMENIIDAE**
Epermenia chaerophylella (Goeze, 1776)
- PTEROPHORIDAE**
Platyptilia gonodactyla ([Denis & Schiffermüller], 1775)
Gillmeria ochrodactyla ([Denis & Schiffermüller], 1775)
Stenoptilia bipunctidactyla (Scopoli, 1763)
Cnaemidophorus rhododactylus ([Denis & Schiffermüller], 1775)
Oxyptilus pilosellae (Zeller, 1841)

- Crombrugghia distans* (Zeller, 1847)
Pterophorus pentadactylus (Linnaeus, 1758)
Hellinsia lienigiana (Zeller, 1852)
Adaina microdactyla (Hübner, 1813)
Emmelina monodactyla (Linnaeus, 1758)
- CARPOSINIDAE**
Carposina scirrhosella Herrich-Schäffer, 1854
- THYRIDIDAE**
Thyris fenestrella (Scopoli, 1763)
- PYRALIDAE**
Aphomia sociella (Linnaeus, 1758)
Aphomia zelleri Joannis, 1932
Galleria mellonella (Linnaeus, 1758)
Synaphe punctalis (Fabricius, 1775)
Pyralis farinalis Linnaeus, 1758
Hypsopygia costalis (Fabricius, 1775)
Hypsopygia incarnatalis (Zeller, 1847)
Hypsopygia rubidalis ([Denis & Schiffermüller], 1775)
Hypsopygia glaucinalis (Linnaeus, 1758)
Endoricha flammealis ([Denis & Schiffermüller], 1775)
Cryptoblabes bistriga (Haworth, 1811)
Trachonitis cristella ([Denis & Schiffermüller], 1775)
Sciota rhenella (Zincken, 1818)
Sciota hostilis (Stephens, 1834)
Sciota adelphella (Fischer von Röslerstamm, 1836)
Selagia argyrella ([Denis & Schiffermüller], 1775)
Oncocera semirubela (Scopoli, 1763)
Phycita roborella ([Denis & Schiffermüller], 1775)
Nephopterix angustella (Hübner, 1796)
Acrobasis tumidana ([Denis & Schiffermüller], 1775)
Acrobasis repandana (Fabricius, 1798)
Acrobasis consociella (Hübner, 1813)
Gymnancyla hornigi (Lederer, 1852)
Assara terebrella (Zincken, 1818)
Euzophera fuliginosella (Heinemann, 1865)
Homoeosoma sinuella (Fabricius, 1798)
Homoeosoma nebulella ([Denis & Schiffermüller], 1775)
Phycitodes binaevelia (Hübner, 1813)
Phycitodes inquinatella (Ragonot, 1887)*
Phycitodes albatella (Ragonot, 1887)*
Plodia interpunctella (Hübner, 1813)
Hypsotropa unipunctella Ragonot, 1888
Ematheudes punctella (Treitschke, 1833)
- CRAMBIDAE**
Scoparia pyralella ([Denis & Schiffermüller], 1775)
Scoparia ingratella (Zeller, 1846)
Scoparia ambigualis (Treitschke, 1829)
Scoparia subfusca Haworth, 1811
Scoparia basistrigalis Knaggs, 1866
Eudonia mercurella (Linnaeus, 1758)
Eudonia pallida (Curtis, 1827)
- Euchromius ocellea* (Haworth, 1811)
Chilo phragmitella (Hübner, 1805)
Calamotropha paludella (Hübner, 1824)
Chrysoteucha culmella (Linnaeus, 1758)
Crambus pascuella (Linnaeus, 1758)
Crambus lathoniellus (Zincken, 1817)
Crambus perlella (Scopoli, 1763)
Agriphila deliella (Hübner, 1813)
Agriphila tristella ([Denis & Schiffermüller], 1775)
Agriphila inquinatella ([Denis & Schiffermüller], 1775)
Agriphila straminella ([Denis & Schiffermüller], 1775)
Agriphila tolli pelsonius Fazekas, 1985
Catoptria permutterella (Herrich-Schäffer, 1848)*
Catoptria pinella (Linnaeus, 1758)
Catoptria falsella ([Denis & Schiffermüller], 1775)
Catoptria verella (Zincken, 1817)
Mesocrambus candiellus (Herrich-Schäffer, [1848])
Pediasia luteella ([Denis & Schiffermüller], 1775)
Pediasia contaminella (Hübner, 1796)
Platytes cerusella ([Denis & Schiffermüller], 1775)
Platytes alpinella (Hübner, 1796)
Ancylolomia palpella ([Denis & Schiffermüller], 1775)
Schoenobius gigantella ([Denis & Schiffermüller], 1775)
Donacaula forficella (Thunberg, 1794)
Donacaula mucronella ([Denis & Schiffermüller], 1775)
Elophila nymphaea (Linnaeus, 1758)
Acentria ephemerella ([Denis & Schiffermüller], 1775)
Cataclysta lemnata (Linnaeus, 1758)
Parapoynx stratiotata (Linnaeus, 1758)
Aporodes floralis (Hübner, 1809)
Cynaeda dentalis ([Denis & Schiffermüller], 1775)
Epascestria pustulalis (Hübner, 1823)
Evergestis forficalis (Linnaeus, 1758)
Evergestis extimalis (Scopoli, 1763)
Evergestis limbata (Linnaeus, 1767)
Evergestis pallidata (Hufnagel, 1767)
Paracorsia repandalis ([Denis & Schiffermüller], 1775)
Ecpyrrhorhoe rubiginalis (Hübner, 1796)
Pyrausta despicata (Scopoli, 1763)
Pyrausta aurata (Scopoli, 1763)
Pyrausta purpuralis (Linnaeus, 1758)
Nascia cilialis (Hübner, 1796)
Sitochroa palealis ([Denis & Schiffermüller], 1775)
Sitochroa verticalis (Linnaeus, 1758)
Anania coronata (Hufnagel, 1767)
Anania crocealis (Hübner, 1796)
Anania fuscalis ([Denis & Schiffermüller], 1775)
Anania hortulata (Linnaeus, 1758)
Anania lancealis ([Denis & Schiffermüller], 1775)
Anania perlucidalis (Hübner, 1809)
Anania stachydalis (Zincken, 1821)
Anania verbascalis ([Denis & Schiffermüller], 1775)
Sclerocona acutella (Eversmann, 1842)
Psammotis pulveralis (Hübner, 1796)
Ostrina nubilalis (Hübner, 1796)

| | |
|---------------------------------------------------------|-------------------------------------------------------------|
| <i>Paratalanta pandalis</i> (Hübner, 1825) | <i>Agrotera nemoralis</i> (Scopoli, 1763) |
| <i>Paratalanta hyalinalis</i> (Hübner, 1796) | <i>Diasemia reticularis</i> (Linnaeus, 1761) |
| <i>Udea ferrugalis</i> (Hübner, 1796) | <i>Metasia ophialis</i> (Treitschke, 1829) |
| <i>Pleuroptya ruralis</i> (Scopoli, 1763) | <i>Nomophila noctuella</i> ([Denis & Schiffermüller], 1775) |
| <i>Mecyna flavalis</i> ([Denis & Schiffermüller], 1775) | |

Acknowledgements

The author offers a word of thanks to H. W. Van der Wolf (NL), J. C. Koster (NL), W. Biesenbaum (D) and O. Karsholt (DK) for help with identifying difficult species. Thanks are to G. Pastorális (SK) for the invaluable and advice. Barry Goater (GB-Chandlers Ford) corrected the English language of the manuscript. We are grateful to all for their help.

References

- ÁCS E. & SZABÓKY Cs. 1993: The Lepidoptera fauna of the Bükk National Park. – In: MAHUNKA S. (ed.): The fauna of the Bükk National Park I. Hungarian Natural History Museum, Budapest, 186–220.
- BALDIZZONE, G., VAN DER WOLF, H. W. & LANDRY, J.-F. 2006: Coleophoridae, Coleophorinae (Lepidoptera). In: World Catalogue of Insects 5. – Apollo Books, Stenstrup, 215 pp.
- BIESENBAUM, W. & VAN DER WOLF, H. W. 1999: Die Lepidopterenfauna der Rheinlande und Westfalens, Band 7. Familie: Coleophoridae Hübner [1825]. – Arbeitsgemeinschaft rheinisch-westfälischer Lepidopterologen e. V. Düsseldorf, 333 pp., Taf. I–XXIX.
- BUSCHMANN F. 2003: A Mátra Múzeum molylepke-gyűjteménye I. Micropterigidae – Gelechiidae. – Folia Historico Naturalia Musei Matraensis 27: 267–287.
- BUSCHMANN F. 2004: A Mátra Múzeum molylepke-gyűjteménye III. Choreutidae – Pyralidae. – Folia Historico Naturalia Musei Matraensis 28: 243–272.
- BUSCHMANN F. 2005: Új microlepidoptera fajok a Mátra Múzeum gyűjteményében – Folia Historico Naturalia Musei Matraensis 26: 173–175.
- ELSNER, G., HUEMER, P. & TOKÁR, Z., 1999: Die Palpenmotten (Lepidoptera: Gelechiidae) Mitteleuropas. Bestimmung – Verbreitung – Flugstandort. – František Slamka, Bratislava, 208 pp.
- FARKAS S. 1992: Vizsgálatok Tolna megyei természetvédelmi területek nappali lepke faunáján. – Folia Entomologica Hungarica 53: 253–255.
- FAZEKAS I. 1986: Ergänzungen zur Verbreitung europäischer Crambinae- und Pterophoridae-Arten (Lepidoptera). – Entomologische Zeitschrift 96 (17): 245–253.
- FAZEKAS I. 1991: Cochyli flaviciliiana Westwood, 1854 und Stenoptilia plagiодactyla Stainton, 1851 als neue Arten für Ungarns Fauna. – Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N. F. 12: 202–210.
- FAZEKAS I. 1992a: Tolna megye nappali lepkéi. [Die Tagfalterfauna des Komitats Tolna, Ungarn]. – Babits-füzetek 7: 1–142.
- FAZEKAS I. 1992b: Új Pyralidae fajok Nyugat-Magyarországon. – Savaria 20/2: 49–54.
- FAZEKAS I. 1993: A Tihanyi Tájvédelmi Körzet lepkafaunája (1.). Faunisztkai alapvetés (Lepidoptera). – A Bakonyi Természettudományi Múzeum Közleményei 12: 105–144.
- FAZEKAS I. 1994a: A magyarországi makrorégiók Cochylini faunája (Lepidoptera: Tortricidae) I. A Dunántúli dombság. – Állattani Közlemények 80: 35–56.
- FAZEKAS I. 1994b: Das Cochylini-Material aus Ungarn des Wiener Naturhistorischen Museums und der Zoologischen Staatsammlung München. – Nachrichtenblatt der Bayerischen Entomologen 43: 39–46.
- FAZEKAS I. 1996: Phycitodes inquinatella exustella (Ragonot, 1888) in Hungary, Pyralidae. – Állattani Közlemények 81: 15–17.
- FAZEKAS I. 1998: Daten zur Kenntnis der Pyraloidea-Fauna Ungarns (Nr.1). – Folia Comloensis 7: 49–66.
- FAZEKAS I. 2001a: Somogy megye molylepke faunája (Lepidoptera: Microlepidoptera). – Natura Somogyiensis 1: 303–327.

- FAZEKAS I. 2001b: A Coleophora chrysanthemi O. Hoffmann, 1896 és az Ancylosis roscidella (Eversmann, 1844) új molylepke fajok Magyarországon (Microlepidoptera: Coleophoridae, Pyralidae). – Folia Historico Naturalia Musei Matraensis 25: 253–260.
- FAZEKAS I. 2001c: A Mátra-vidék Pyraloidea (s. str.) faunája (Microlepidoptera). – Folia Historico Naturalia Musei Matraensis 25: 261–286.
- FAZEKAS I. 2002a: Systematisches und synonymisches Verzeichnis der Microlepidoptera Ungarns (Lepidoptera: Microlepidoptera). – Folia Historico Naturalia Musei Matraensis 26: 289–327.
- FAZEKAS I. 2002b: Baranya megye Microlepidoptera faunájának katalógusa (Lepidoptera). – Folia Comloensis 11: 5–76.
- FAZEKAS I. 2008: A Scythris sinensis Felder & Rogenhofer, 1875 új lelőhelye Magyarországon (Microlepidoptera: Scythrididae). [New record of the Scythris sinensis Felder & Rogenhofer, 1875 in Hungary (Microlepidoptera: Scythrididae)]. – Acta Naturalia Pannonica 3, Suppl. 2: 169–172.
- FAZEKAS I. 2008a: Microlepidoptera Pannoniae meridionalis, VII. Faunisztikai és taxonómiai adatok Somogy megyéből (1.) (Lepidoptera). – Somogyi Múzeumok Közleményei 18: 101–115.
- FAZEKAS I. 2008b: A Scythris sinensis Felder & Rogenhofer, 1875 új lelőhelye Magyarországon (Microlepidoptera: Scythrididae). New record of the Scythris sinensis Felder & Rogenhofer, 1875 in Hungary (Microlepidoptera: Scythrididae). – Acta Naturalia Pannonica 3, Suppl. 2: 169–172.
- FAZEKAS I. 2010: Coleophora alnifoliae Barasch, 1934 and Alucita palodactyla Zeller, 1847 in Hungary (Lepidoptera: Coleophoridae & Alucitidae). – e-Acta Naturalia Pannonica, 1 (2): 205–212.
- GOZMÁNY L. 1955: Molylepkék III. Microlepidoptera III. – Fauna Hungariae XVI., 4: 64 pp.
- GOZMÁNY L. 1956: Molylepkék II. Microlepidoptera II. – Fauna Hungariae XVI., 3: 136 pp.
- GOZMÁNY L. 1958: Molylepkék IV. Microlepidoptera IV. – Fauna Hungariae XVI., 5: 295 pp.
- GOZMÁNY L. 1963: Molylepkék VI. Microlepidoptera VI. – Fauna Hungariae XVI., 7: 289 pp.
- GOZMÁNY L. 1968: Hazai molylepkéink magyar nevei. – Folia Entomologica Hungarica 21: 225–296.
- GOZMÁNY L. & SZÖCS J. 1965: Molylepkék I. Microlepidoptera I. – Fauna Hungariae XVI., 2: 214 pp.
- KARSHOLT, O. & NIEUKERKEN, E. J. VAN (eds.) 2005. Lepidoptera, Moths. Fauna Europaea, version 1.2, <http://www.faunaeur.org>
- KIMBER, I. 2010: Oecophoridae, 644 Borhausenia fuscescens (Haworth, 1828). – <http://ukmoths.org.uk> [visited 21.04.2010].
- KOSTER, J. C. & SINEV, S. YU. 2003: Momphidae, Batrachedridae, Stathmopodidae, Agonoxenidae, Cosmopterigidae, Chrysopidae. – In HUEMER, P., KARSHOLT, O. & LYNEBORG (eds): Microlepidoptera of Europe 5: 1–387.
- PAP N. 2007: Tolna – a rural area in Central-Europe Regional and local development in Tolna County, Hungary. – Lomart Kiadó, 164 pp.
- PASTORÁLIS G. 2010: A checklist of Microlepidoptera (Lepidoptera) occurred in the territory of Hungary (1.4). – e-Acta Naturalia Pannonica 1 (1): 89–170.
- PILlich F. 1914: Aus der Arthropodenwelt Simontornya's. – Simontornya, Hungaria occidentalis, Komitat Tolna, 171 pp.
- RONKAY L. & SZABÓKY Cs. 1981: Investigations on the Lepidoptera fauna of the Zemplén Mts. (NE Hungary). I. The valley of Kemence stream. – Folia Entomologica Hungarica 42: 167–184.
- SZABÓKY Cs. 1983: A Dé-Dunántúl molylepkéi I. Nattán Miklós molylepke-gyűjteménye (Lepidoptera). – A Janus Pannonius Múzeum Évkönyve 27: 15–35.
- SZABÓKY Cs. 1994: Adatok a magyar faunára új molylepkéről. – Folia Entomologica Hungarica 55: 381–383.
- SZABÓKY Cs. 1998: Molyfaunisztikai újdonságok III. – Folia Entomologica Hungarica 59: 305–308.
- SZABÓKY Cs. 2004: Molyfaunisztikai újdonságok VII. (Lepidoptera: Coleophoridae, Elachistidae, Gelechiidae, Tortricidae). – Folia Entomologica Hungarica 65: 248–252.
- SZABÓKY Cs. 2009: Pécsely lepkéi (Lepidoptera). – Folia Musei Historico-Naturalis Bakonyiensis 26: 111–140.
- SZEŐKE K. 2009: Gyűrűfű molylepkéin (Microlepidoptera) végzett biodiverzitás vizsgálat eredményei. – Natura Somogyiensis 13: 163–168.
- SZIRÁKY Gy. 1980: Notes on Coleophora and Cnephasia species trapped by synthetic attractans (Lepidoptera: Coleophoridae and Tortricidae). – Folia Entomologica Hungarica 33 (1): 161–166.
- SZÖCS J. 1973: Újabb molylepkék a magyar faunában. – Folia Entomologica Hungarica, 26: 155–169.
- SZÖCS J. 1977a: A lepkéhez kötött természetes tápnövényei, III. – Folia Entomologica Hungarica 30: 143–150.
- SZÖCS J. 1977b: Lepidoptera – aknák és gubacsok – Fauna Hungariae XVI. 16: 423 pp.
- TAKÁCS A. 2009: Gánt-Gránás lepkéi (Lepidoptera). – Folia Musei Historico-Naturalis Bakonyiensis 26: 141–170.